

### REMARKS/ARGUMENTS

Reconsideration of this application is requested. Claims 1-6, 8, 10 and 14-18 are pending in the application subsequent to entry of this Amendment.

The claims have been amended in order to more particularly point out and distinctly claim that which applicants regard as their invention, to address claim clarity issues raised on pages 4-5 of the Official Action, to reduce issues and to advance prosecution generally.

Claims directed to compounds *per se* have been revised to refer to a single compound and adjusted to include proper terminology, as appropriate. Claims 7 and 9 have been canceled to reduce issues, claim 10 revised to depend from claim 8. Claims 11-13 have been canceled and replaced with new method claim 15-18. New claim 14 has been added directed to the specific amino acids listed in the first paragraph of page 7 of the description. Further comments regarding the claim amendments will appear in the discussion that follows.

#### Claims 9-13 – 35 USC § 112

Claims 9 and 11-13 are canceled and replaced with new method of treatment claims 14-18. Claim 10 is dependent from claim 8 directed to a pharmaceutical composition; claim 10 should not have been included in this rejection. This response is made on the basis of new claims 14-18.

Camptothecins are a well-known class of antitumor drugs used for the treatment of different tumors. The description of the invention makes explicit reference to EP 1 044 977 (EP '977) and the literature cited therein. This reference reports that camptothecins demonstrated a wide spectrum of antitumor activity, in particular against colon tumors, other solid tumors and leukemias (see page 1, fourth paragraph). The clinical reference for camptothecins is Topotecan, which is indicated for the treatment of small and non-small cell lung, ovarian, breast, stomach, liver, prostate, soft tissue sarcoma, head and neck tumors, cancer of the esophagus, resistant colon-rectum, multiform glioblastoma, and chronic and acute myelocytic leukemias (EP'977, page 3, third paragraph). The compounds disclosed in EP'977, of which the compounds of the present invention are derivatives as far as the 7-C(R<sub>5</sub>)=N-O-R<sub>4</sub> substituent is concerned, and the activity on a wide range of tumors is proven by the enclosed list of references retrieved from PubMed (Annex 4). See also Penco et al., US 6.242.457, claiming priority of EP'744, where the range of treated tumors was expanded (note column 14).

BEST AVAILABLE COPY

Further evidence of a wide range of antitumor activity is found in the same references as cited by the examiner, (*see* WO'691).

Therefore, the state of the art as to the therapeutic benefits to be obtained from the camptothecins must be recognized, since there is no reason for one skilled in this art to believe that the claimed compounds would not be effective on different tumors. To the contrary, the problem solved by the present invention is to enhance stability and solubility in water of the molecule, thus enhancing the pharmacological activity and giving a person skilled in the art a more powerful therapeutic tool in terms of an improved therapeutic index; *see* page 2, last line.

The present compounds are not claimed for treating tumors which are not usually treated with Topoisomerase I inhibitors or with camptothecins in general. This is very clear from the entire content of the description and the references cited in it and is clearly stated in new claims 15-17. Therefore, the skilled clinician will know for which type of susceptible tumors the compounds can be used.

The activity of camptothecins in parasitic and viral infections is also well-known in the art (*see* WO'876, page 23, first full paragraph) and thus is claimed in new claim 18.

In view of the above, applicants respectfully request withdrawal of this rejection.

Claims 1, 4-6, 7 and 9-13 – 35 USC § 112

Claim 1: there are only two cases for n and m:

- 1) both n and m are 1;
- 2) both n and m are 0.

No other cases are disclosed and this is clarified in amended claim 1.

The definition of residue B as amino acid is clearly explained on page 7, first paragraph of the description and claim 1 is amended to reflect this description. New dependent claim 14 is added directed to the amino acids specified in this passage.

R<sub>4</sub> seems clearly defined by the list of variables in claim 1. The term "residue" has been replaced with "substituent".

The spelling of "benzylglycil" has been corrected in claim 4.

Claim 5: actually, step b) as written is self-explanatory. Substitution of the leaving group with the Y group is a well-known class of chemical reactions, also known as nucleophilic substitutions. Please see the attached chapter of a university textbook (Annex 1) explaining the

mechanism of nucleophilic substitution reactions. The term "substitution of said leaving group with the Y group" recited in claim 5 is sufficient to indicate to the skilled person what to do to carry out the claimed process to obtain the desired product. In any event, all one has to do is look in applicants' specification. Claim 5 is supported by the description, see page 10, "preparation of the intermediate products 2a,b" and page 11, "preparation of the intermediate products 4a,b". Claim 5 will be interpreted in light of the description.

Claim 6: also step b) is self-explanatory as written. Transformation of a carboxylic group into an amide is a well-known chemical reaction. Please see the attached copy of a chapter of a university textbook (Annex 2) showing different modes for obtaining an amide. In any event, all one has to do is look in applicants' specification. Claim 6 is supported by the description, see page 11, "preparation of the intermediate products 6a,b" and page 12, "preparation of the intermediate products 7a,b" and "preparation of the products 8a,b". Claim 6 will be interpreted in light of the description.

Claim 7 has been deleted.

Claim 9 has been deleted.

Claim 10, a pharmaceutical composition claim dependent from claim 8, is maintained. Applicants believe this claim is clear to the skilled reader, since combining different antitumor drugs is common practice in this field. The other antitumor drug will be determined by the skilled person simply exercising his or her knowledge. Attached are only the only first 50 titles out of 269 reviews published until 2002 (priority year of this application) to show the current knowledge of cancer polychemotherapy (Annex 3).

The subject matter of claims 12 and 13 is maintained and written as methods of treatments in new claims 14-17, since applicants believe that the skilled person will know how to use the invention just resorting to common knowledge. A sample of literature is enclosed (Annex 5). The examiner has already indicated the treatment of lung tumor is enabled; *see* item 4, second line and compare with new claim 17. Claim 16 is directed to a particular group of susceptible tumors described in the paragraph bridging pages 13-14 of the specification.

Claims 1, 3, 5, 7-9 and 12 – 35 USC § 102(b)

Claim 1 has been amended by deleting the meaning of hydrogen for the R<sub>1</sub> group. This amendment restores novelty over Vishnuvajjala.

Claims 1, 2, 4 and 6-13 – 35 USC § 103(a)

The Examiner admitted that the prodrug disclosed by Matsumoto et al. is an HIV protease inhibitor, but is of the opinion that the document affirms the general applicability of the approach and points at Figure 2 to support the opinion.

The Matsumoto et al. reference is cited in the subject application (page 2) with the following comment: *"molecule of a dipeptide nature, differing enormously from the molecular structure of camptothecin, and to that end functionalise a hydroxyl group with a portion formed by a spacer part and a solubilizing part. The spacer part is provided by a bicarboxylic acid, whereas the solubilizing part is provided by a diamine"*.

Matsumoto et al. give only more than general comments on the possible solubilizing effect of the system provided by them, namely the spacer and the solubilizing portion, the latter releasing the drug through a cyclization mechanism.

The skilled person must find in the prior art a clear teaching of the case this person is facing, with all instructions and indication on how to obtain the expected results. In the present case, Matsumoto et al. with all its insufficiencies does not apply, neither alone nor in combination with Penco et al.

Penco et al. mention the problem of water-solubility of camptothecin derivatives, but give no indication on how to solve this problem.

Matsumoto et al. enable only for a dipeptide derivative. Therefore, this reference is not predicable for camptothecin derivatives, which are polycyclic compounds of a completely different chemical nature.

Secondly, Matsumoto et al. put very strict rules in designing the prodrug: a) only succinic or glutaric dicarboxylic acids are provided, and a short chain amine as solubilizing portion are disclosed. The final teaching of Matsumoto et al. is to find suitable conditions to enhance cyclization of the spacer-solubilizing system, and this leads to short chain elements ending with an amino group. The authors advise against different ending groups (see page 608, right column).

Ultimately, Matsumoto et al. teach a solubilizing system which will cyclize and release the drug. There is the least suggestion that such a system with strict rules will work with 7-substituted camptothecins derivatives, as the claimed ones.

Moreover, the present invention goes against Matsumoto et al. teaching, by providing a wide range of chain length (from 1 to 8 carbon atoms) and a wide variety of terminal groups other than a simple amino group.


Therefore, Matsumoto et al. give no expectation to successfully solve the problem of water-solubility passing from a dipeptide derivative, such as the only enabled compound KNI-727, to a complex polycyclic compound, bearing a complex substitution in position 7, such as the camptothecin derivatives of the present invention.

For the above reasons it is respectfully submitted that the claims of this application are in proper order, enabled and define subject that is patentable over the cited and applied prior art references. Reconsideration and favorable action are solicited. Should the examiner have any questions or require further information, please contact the undersigned.

Respectfully submitted,

**NIXON & VANDERHYE P.C.**

By: \_\_\_\_\_

  
Arthur R. Crawford  
Reg. No. 25,327

ARC:eaw  
901 North Glebe Road, 11th Floor  
Arlington, VA 22203-1808  
Telephone: (703) 816-4000  
Facsimile: (703) 816-4100

*A Series of Books in Organic Chemistry*  
Andrew Streitwieser, Jr., Editor

ANNEX 1

# ***INTRODUCTION 1***

# ***ORGANIC CHEMISTRY***

***Andrew Streitwieser, Jr.***

***Clayton H. Heathcock***

*University of California, Berkeley*

***Macmillan Publishing Co., Inc.***

***NEW YORK***

***Collier Macmillan Publishers***

***LONDON***


## CHAPTER 8

### Reactions of Alkyl Halides


8.1

#### The Displacement Reaction

The replacement of the halogen in an alkyl halide by another group is one of the most important reactions in organic chemistry. In Section 3.1, we took a brief look at one such reaction, the reaction of methyl chloride with hydroxide ion.



Another example is the reaction of ethyl bromide with potassium iodide in acetone solution.



Although this is an equilibrium process, the reaction proceeds virtually to completion because potassium iodide is soluble in acetone and potassium bromide is not.

Like the reaction discussed in Section 3.1, the reaction of ethyl bromide with iodide ion is relatively slow. In order for complete reaction to occur, it is necessary to heat the mixture for several hours. The rate of the reaction may be determined by following the rate of disappearance of reactants or the rate of appearance of products. It is proportional to the product of the concentrations of the two reactants.

$$\text{rate} = -\frac{d[\text{C}_2\text{H}_5\text{Br}]}{dt} = -\frac{d[\text{I}^-]}{dt} = \frac{d[\text{C}_2\text{H}_5\text{I}]}{dt} = \frac{d[\text{KBr}]}{dt} = k[\text{C}_2\text{H}_5\text{Br}][\text{I}^-]$$

This equation is expressed in the symbolism of calculus. The expression

$$\frac{d[\text{C}_2\text{H}_5\text{Br}]}{dt}$$

means simply the rate with which the concentration of  $\text{C}_2\text{H}_5\text{Br}$  changes with time. The negative sign indicates that the concentration of  $\text{C}_2\text{H}_5\text{Br}$  decreases as time increases. Note that in this case  $[\text{KBr}]$  refers to amount rather than concentration because of its low solubility in acetone.

The concentrations of  $\text{C}_2\text{H}_5\text{Br}$ ,  $\text{I}^-$ ,  $\text{C}_2\text{H}_5\text{I}$ , and  $\text{KBr}$  may be determined at different times during the reaction by chemical or spectroscopic analysis. As the reaction proceeds, the concentrations of the reactants become reduced and the rate of reaction decreases. For example, at  $50^\circ$ , with the two reactants each present in an initial concentration of  $0.1\text{ M}$ , the reaction is 50% complete in 7 min but only 95% complete after 2 hr. Furthermore, the reaction has an activation energy,  $\Delta H^\ddagger$ , of  $19\text{ kcal mole}^{-1}$ ; it is 10 times slower at  $25^\circ$  than it is at  $50^\circ$ . This activation energy is considerably higher than those of the free radical reactions we studied in Chapter 5.

When the rate of a chemical reaction depends on the concentration of two



$$I^- + \begin{array}{c} Br \\ | \\ CH_3-C-H \\ | \quad \backslash \\ H \quad H \end{array} \longrightarrow \begin{array}{c} I \\ | \\ CH_3-C-H \\ | \quad \backslash \\ H \quad H \end{array} + Br^-$$

This does not mean that this is not a perfectly good mechanism, only that another mechanism is better for this particular reaction. The activation energy of the frontal-attack mechanism is so much higher than that of the actual mechanism that no significant number of product molecules are formed by such a path. Since rates are exponential functions of activation energy, a small energy change can have a dramatic effect on rate. For example, if two reactions differ in activation energy by 10 kcal mole<sup>-1</sup>, the ratio of their reaction rates is 10,000,000:1. Thus, it is only necessary that we consider the most probable reaction mechanisms for a given reaction—those with the lowest activation energies.

-82

It has long been known that, when an optically active alkyl halide is exposed to halide ion in solution, the optical activity gradually diminishes to zero. This is an example of racemization; the optically active halide is converted to an equimolar mixture of (+) and (-) enantiomers. The rate of racemization is dependent on the concentration of both the alkyl halide and the added halide. The rate constant for the racemization process is called  $k_a$ . The conditions under which such racemization occurs are the same as those that lead to substitutions such as those discussed in Section 8.1.

frontal attack mechanism:

$$\begin{array}{c} \text{CH}_3 \\ | \\ \text{C} - \text{H} \\ / \quad \backslash \\ \text{C}_2\text{H}_5 \quad \text{I} \\ (\text{R}) \end{array} + \text{X}^- = \begin{array}{c} \text{CH}_3 \\ | \\ \text{C} - \text{H} \\ / \quad \backslash \\ \text{C}_2\text{H}_5 \quad \text{X} \\ (\text{R}) \end{array} + \text{I}^-$$

There must be a mechanism for substitution that allows racemization at C-2. To

## Sec. 8.2


Stereochemistry  
of the  
Displacement  
Reaction

determine how often each replacement of one halide by another is accompanied by a change from the (+) enantiomer to the (-) enantiomer, it is necessary to know both the rate of racemization and the rate of the substitution reaction itself.


This experiment was first done in 1935 by the late Professor E. D. Hughes and his colleagues at the University of London. The experiment was an extremely elegant one, which used sodium iodide containing a radioactive isotope of iodine,  $^{128}\text{I}$ , as the attacking group and optically active 2-iodooctane containing normal  $^{127}\text{I}$  as the alkyl halide.

$^{128}\text{I}$  is prepared by exposing normal  $^{127}\text{I}$  to neutrons. The radioisotope decays with a half-life of 25 min.


At equilibrium, the radioactive iodine is distributed equally between 2-iodooctane and iodide ion.



The reaction was found to have a rate constant at  $30^\circ$ ,  $k_{\text{exch}} = (1.36 \pm 0.11) \times 10^{-3} \text{ M}^{-1} \text{ sec}^{-1}$ . Under identical conditions, the rate constant for racemization of optically active 2-iodooctane was found to be  $k_{\text{a}} = (2.62 \pm 0.03) \times 10^{-3} \text{ M}^{-1} \text{ sec}^{-1}$ . Within experimental error, the rate of racemization is exactly twice that of exchange. Consequently, each act of replacement of one iodide by the other is accompanied by a change from one enantiomer to another. In this way, the rotation of one molecule of product cancels that of one reactant molecule, and is equivalent to two molecules of racemic material. Looking at it in a slightly different way, all of the beginning molecules must undergo reaction with  $^{128}\text{I}^-$  before incorporation of the radioisotope is complete, whereas only one half of the beginning molecules must undergo inversion of stereochemistry before racemization is complete. Thus, if inversion of stereochemistry occurs every time substitution occurs,  $k_{\text{a}}$  must be exactly twice  $k_{\text{exch}}$ , as is found.



This observation has been combined with many other studies to infer that, in a bimolecular displacement reaction, the attacking group attacks at the carbon atom at the rear of the bond to the leaving group. During reaction, the carbon forms a progressively stronger bond to the attacking group, while the bond to the leaving group is being weakened. During this change, the other three bonds to the central carbon progressively flatten out and end up on the other side of the carbon in a manner similar to the spokes of an umbrella inverting in a windstorm.



The reaction is formulated as above, with the transition state geometry in brackets. The dotted lines indicate a partially formed or partially broken bond. The symbol

## Chap. 8

Reactions of  
Alkyl Halides

$\delta^-$  indicates that the negative charge is spread over both iodines in the transition state.

During the course of the reaction, the reacting system has greater potential energy than either the reactants or the products. The two weak bonds entering and leaving groups are weaker than the single bond in either the reactant or the product. Hence, energy is required in order for reaction to occur. The necessary potential energy is supplied by the conversion of kinetic energy; only the minute fraction of reactants that have sufficient kinetic energy can undergo reaction. Furthermore, even if the colliding reactants have sufficient kinetic energy, they must have the proper orientation or they will simply bounce apart. The point of highest energy is called the **transition state**. It is important to remember that the transition state is a point of *maximum* energy. It is not a molecule that can be isolated and studied. In fact the whole act of displacement occurs in the space of about  $10^{-12}$  sec, the period of a single vibration. The reacting system has the transition state geometry for only a fleeting moment.

The geometry of the transition state appears to be that in which the entering and leaving groups are both weakly bonded to carbon in a linear fashion in which the three remaining bonds to carbon lie in a plane perpendicular to the two weak bonds. The reaction mechanism for reaction of an entering nucleophile,  $Y^-$ , and a leaving group,  $X^-$ , is shown in Figure 8.1, where the structure of the reacting system at several points along the reaction coordinate is illustrated. At point (b) the  $C-X$  bond has started to lengthen and the central carbon has started to flatten out. At the transition state, point (c), the central carbon is approximately flat and both the bonds to the leaving and entering group are long. At point (d) the  $C-Y$  bond has started to lengthen and the central carbon has started to flatten out.




FIGURE 8.1 Reaction mechanism profile for a displacement reaction by  $Y^-$  on  $RX$ .

## Sec. 8.2

Stereochemistry  
of the  
Displacement  
Reaction

occurs on the final road to products (e); the central carbon has bent, the C—Y bond is approaching normal length, and the leaving group, X, is receding. The structures at points (a) through (e) are represented in stereo form in Figure 8.2.




FIGURE 8.2 The structure of the reaction system at points (a) through (e) in Figure 8.1.

In orbital terms, both the reactant and the product are tetrahedral. The C—X bond in each case is  $C_{sp^3}$ -X. In the transition state, the weak bonds to X and Y may be considered to derive from overlap of a halogen orbital with the two lobes of a  $p$  orbital on the central carbon. The other three bonds to this carbon are formed from  $sp^2$  hybrid orbitals, as shown in Figure 8.3.





FIGURE 8.3 Orbital formulation of the transition state of a displacement reaction.

### 8.3 Generality of the Displacement Reaction

The importance of the displacement reaction lies in its generality. Although the reaction was introduced in the preceding sections with halide ions as entering groups, analogous reactions are known with a wide range of anions and neutral molecules. The only requirement is that the attacking group be a Lewis base, a species which contains an atom that has a pair of electrons available for bonding. The examples in Table 8.1 demonstrate the range of attacking groups that undergo the reaction.

A base may generally be regarded as the conjugate base of an acid, and may be neutral or charged.



A quantitative measure of the basicity of a base is the acidity, or  $pK_a$ ,\* of its conjugate acid.

$$K_a = \frac{[\text{B}^-][\text{H}^+]}{[\text{BH}]} \quad \text{or} \quad \frac{[\text{B}][\text{H}^+]}{[\text{BH}^+]}$$

$$pK_a = -\log K_a$$

A weak conjugate acid (more positive  $pK_a$ ) corresponds to a strong base; a strong

TABLE 8.1  
Some Displacement Reactions with Ethyl Bromide

Attacking Reagent		Product	
Formula	Name	Formula	Name
$\text{HO}^-$	hydroxide ion	$\text{C}_2\text{H}_5\text{OH}$	ethyl alcohol
$\text{C}_2\text{H}_5\text{O}^-$	ethoxide ion	$\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$	diethyl ether
$\text{HS}^-$	hydrosulfide ion	$\text{CH}_3\text{CH}_2\text{SH}$	ethanethiol
$\text{SCN}^-$	thiocyanate ion	$\text{CH}_3\text{CH}_2\text{SCN}$	ethyl thiocyanate
$\text{CN}^-$	cyanide ion	$\text{CH}_3\text{CH}_2\text{CN}$	ethyl cyanide, propionitrile
$\text{N}_3^-$	azide ion	$\text{CH}_3\text{CH}_2\text{N}_3$	ethyl azide
$\text{NH}_3$	ammonia	$\text{CH}_3\text{CH}_2\text{NH}_3^+ \text{Br}^-$	ethylammonium bromide
$\text{H}_2\text{O}$	water	$\text{CH}_3\text{CH}_2\text{OH}_2^+ \text{Br}^-$	ethyloxonium bromide
$\text{CH}_3\text{CO}_2^-$	acetate ion	$\text{CH}_3\text{CO}_2\text{C}_2\text{H}_5$	ethyl acetate
$\text{NO}_3^-$	nitrate ion	$\text{CH}_3\text{CH}_2\text{ONO}_2$	ethyl nitrate
$\text{P}(\text{CH}_3)_3$	trimethylphosphine	$\text{C}_2\text{H}_5\text{P}(\text{CH}_3)_3^+ \text{Br}^-$	ethyltrimethyl- phosphonium bromide
$\text{N}(\text{C}_2\text{H}_5)_3$	triethylaniline	$(\text{C}_2\text{H}_5)_4\text{N}^+ \text{Br}^-$	tetraethylammonium bromide

\* For a review of  $pK$ , see sections 11.4, 17.4, and Appendix IV.

## Sec. 8.3

Generality of the  
Displacement  
Reaction

conjugate acid (less positive  $pK_a$ ) corresponds to a weak base. A few common bases, with the  $pK_a$ s of the corresponding conjugate acids, are given in Table 8.2. Other values are given in Appendix IV.

The basicity of a base depends on the strength of its bond to a proton in solution. In the displacement reaction, the base forms a bond to carbon. Consequently, we might expect to find some correlation between the energy involved in the protonation of a base and the activation energy for its reaction in a displacement reaction. Instead of energies or enthalpies, many correlations of this type have been found using the Gibbs free energies,  $\Delta G$ . The  $pK$  of an acid is proportional to the standard free energy of the acid-base equilibrium:

$$\Delta G^\circ = 2.303 RT pK$$

This equation follows by combining the relation between an equilibrium constant and the standard free energy with the definition of  $pK$ .

$$\Delta G^\circ = -RT \ln K$$

Recall from Chapter 3 that rate constants are also related to free energies of activation,  $\Delta G^\ddagger$ :

$$k = \text{constant} \times e^{-\Delta G^\ddagger/RT}$$

$$-\log k = \frac{\Delta G^\ddagger}{2.303 RT} - \log (\text{constant})$$

To test the existence of a correlation between basicity and reactivity in displacement reactions, we first need to determine the second-order rate constants for reaction of a series of bases with some common substrate under the same experimental conditions; that is, for the same solvent and temperature. Next, we plot the logarithms of these rate constants against the corresponding  $pK_a$  values. For a perfect correlation every point will fall exactly on a straight line. Figure 8.4 shows a plot of this type for the reactions of a series of bases with methyl iodide in methyl alcohol solution. We see that there is a rough correlation in the anticipated direction. There is a tendency for bases with high  $pK_a$  (stronger bases) to react faster with methyl iodide. The points cluster in a linear fashion but there is a good deal of scatter.

TABLE 8.2  
Some Common Bases

Base	Conjugate Acid	$pK_a$
$I^-$	HI	-9.5
$Br^-$	HBr	-9
$Cl^-$	HCl	-7
$HSO_4^-$	$H_2SO_4$	-5
$H_2O$	$H_3O^+$	-1.7
$F^-$	HF	3.2
$CH_3COO^-$	$CH_3COOH$	4.8
$HS^-$	$H_2S$	7.0
$CN^-$	HCN	9.2
$HO^-$	$H_2O$	15.7
$H_2N^-$	$H_3N$	35

## Chap. 8


Reactions of  
Alkyl Halides

FIGURE 8.4 Correlation of basicity with reactivity toward methyl iodide.

There are a number of reasons why the correlation is not perfect. One is that we have compared reactions in two different solvents, a displacement reaction in methyl alcohol and an acid-base equilibrium in water. Water is a useful solvent for displacement reactions because most organic compounds are not sufficiently soluble. However, only a limited number of  $pK_a$  values are available for nonaqueous solvents.

Another reason why the correlation is not perfect is to be found in the types of bonding in the two systems. In an acid-base equilibrium, the base is involved in a small and relatively "tight"  $1s$  orbital of a hydrogen (Figure 8.5a). In the transition state of a displacement reaction, the orbital containing the lone pair of the base overlaps instead with the larger, more diffuse,  $2p$  orbital of the central carbon (Figure 8.5b).

These two kinds of reactivity of a base are given different names. **Basicity** is the affinity of a base for a proton, and is measured by the equilibrium constant for the reaction with the conjugate acid in water. **Nucleophilicity** (from *nucleus*, L., kernel, and Gr., loving; hence, "nucleus-loving") is the affinity of a base for a carbon atom.




FIGURE 8.5 Comparison of orbital overlaps of a base with a proton and with the central carbon of a displacement reaction.


## Sec. 8.4

## Nucleophilicity

in a displacement reaction transition state. Nucleophilicity is measured by the rate of reaction of the base with a suitable compound, usually methyl bromide or iodide, in some standard solvent.

The displacement reaction is often referred to as an  $S_N2$  reaction for substitution, nucleophilic, bimolecular.

The molecularity of a reaction is defined as the number of molecules involved in the rate-determining transition state. It is sometimes, but not always, equal to the kinetic order of the reaction. For example, consider a reaction of the type



The rate of appearance of the product D is given by the rate law

$$\text{rate of formation of D} = \frac{d[D]}{dt} = \frac{k_1 k_2}{k_{-1}} [A][B] = k' [A][B]$$

which shows second-order kinetics. However, in the slow step ( $k_2$ ), the rate-determining step, only one molecule, species C, is involved. Hence, the reaction is unimolecular.

The mechanistic label  $S_N2$  covers a wide variety of specific reactions (see, for example, Table 8.1 and Figure 8.4). All of the reactions in this category occur by the common mechanism discussed in this section. They all proceed with inversion of configuration at the reacting carbon, and they all show second-order kinetics. The rates of  $S_N2$  reactions are markedly affected by a number of factors, including the nucleophilicity of the attacking group, the structural environment of the carbon where displacement occurs, the nature of the leaving group, and the nature of the solvent. The important principles that enter into evaluating the effect of these variables recur frequently in organic chemistry, and therefore warrant careful study at this time.

## 8.4

## Nucleophilicity

In constructing Figure 8.4, only a fraction of the available data was used. In fact, the bases, or nucleophiles, used in that plot all involve reaction with atoms in the first row of the periodic table—C, N, O, F. The rough correlation evident in Figure 8.4 shows that more basic electron pairs tend to be more nucleophilic.

Now let us add more data and expand the plot to that shown in Figure 8.6. The points in Figure 8.4 have been retained as unlabeled open circles. The additional points for other nucleophilic reagents produce still more scatter. However, a second important generalization can be drawn from Figure 8.6. Second- and third-row elements are invariably more nucleophilic than first-row elements of comparable basicity. The reason for this generalization appears to be that the larger elements have relatively diffuse lone pairs that are more polarizable. These more diffuse lone pairs tend to bond more strongly to the more diffuse  $p$  orbital of the  $S_N2$  transition state than to the small, tight  $1s$  orbital of a hydrogen. Consequently, such lone pairs tend to be more nucleophilic than their basicity would indicate. The first-row elements have smaller orbitals and the lone pairs are held more tightly. These elements are relatively more basic and less nucleo-








FIGURE 8.6 Comparison of nucleophilicities and basicities of various reactants. Open circles refer to points in Figure 8.4.

philic. Examples of reactivity towards methyl iodide are



In each case, the group with the first-row element is the more basic and is less nucleophilic.


At this point we can now interpret an interesting experimental result. The reaction of a sulfate ion is straightforward in its reaction with either H<sup>+</sup> or CH<sub>3</sub>I in an S<sub>N</sub>2 reaction. Both types of reaction occur at an oxygen




## Sec. 8.4

## Nucleophilicity


Sulfite ion, however, behaves quite differently. It reacts with a proton on oxygen to form bisulfite ion, and with methyl iodide on sulfur to form the methanesulfonate ion.




The oxygen in sulfite ion is the more basic atom and prefers to attack  $\text{H}^+$ , but the lone pair on sulfur is more nucleophilic, and it has preference in the  $\text{S}_\text{N}2$  transition state. Sulfate ion has no lone pair on sulfur, and both reactions have no alternative but to occur at the oxygen. Thiosulfate ion is a simple sulfur analog of sulfate. This ion reacts with methyl iodide exclusively on sulfur, even though there are three oxygens and only one sulfur.



Finally, there are some nucleophiles that show measurable nucleophilic properties at two different atoms. Nitrite ion is an example. The ion undergoes protonation exclusively on oxygen to give nitrous acid.




However, the reaction of nitrite ion with methyl iodide gives both methyl nitrite and nitromethane.



In this case, both nitrogen and oxygen are first-row elements and have comparable nucleophilicities. The ratio of the products actually depends on the reaction conditions. Another example is the reaction with cyanide ion. In addition to methyl cyanide, the major product, small amounts of methyl isocyanide are also produced.

**Chap. 8**  
Reactions of  
Alkyl Halides



Anions such as these, which can react at two different positions, are **ambident** (*amho*, L., both; *denris*, L., tooth). "two-fanged" nucleophiles.

### 8.5

#### Effect of Substrate Structure on Displacement Reactions

A large variety of alkyl halides undergo substitution by the  $S_N2$  mechanism. The ease of reaction depends markedly upon the structure of the alkyl to which the halogen is attached. Reactivities vary widely and in a consistent manner. Branching of the chain at the carbon where substitution occurs ( $\alpha$ -carbon) has a significant effect on the rate of reaction. Relative rates of reactions for methyl, ethyl, isopropyl, and *t*-butyl halides are approximately shown in Table 8.3.

**TABLE 8.3**  
*Effect of Branching at the  $\alpha$ -Carbon  
on the Rate of  $S_N2$  Reactions*


Alkyl Halide	Relative Rate
$\text{CH}_3\text{--}$	30
$\text{CH}_3\text{CH}_2\text{--}$	1
$(\text{CH}_3)_2\text{CH--}$	0.03
$(\text{CH}_3)_3\text{C--}$	$\approx 0$

These effects on reaction rate are interpreted with the concept of **steric hindrance** to attack of the attacking nucleophile. The rear of a methyl group is relatively exposed to such attack. As the hydrogens of the methyl group are replaced by methyl groups, the area in the rear of the leaving group becomes more encumbered. It becomes more difficult for the attacking group to approach close enough to the rear of the C—X bond for reaction to occur, and the rate of reaction diminishes (Figure 8.7).



A similar effect may be seen in branching at the  $\beta$ -carbon. Some typical relative rates are shown in Table 8.4. This reduction in rate is also attributable to steric hindrance. In one conformation, the rear of a *n*-propyl carbon is seriously blocked (Figure 8.8a) but in two other conformations, the situation is no worse than for ethyl (Figure 8.8b). Consequently, *n*-propyl halides undergo  $S_N2$  displacement only slightly less readily than do ethyl halides.

For the isobutyl group, it is possible to rotate both of the  $\beta$ -methyl groups out of the way of the attacking group, but the resulting conformation is congested and has relatively high energy (Figure 8.9). Accordingly, isobutyl halides are much less reactive than either ethyl or *n*-propyl compounds.


## Sec. 8.5

Effect of  
Substrate  
Structure on  
Displacement  
ReactionsFIGURE 8.7 Effect of  $\alpha$  branching on  $S_N2$  reactions.TABLE 8.4  
Effect of Branching at the  $\beta$ -Carbon  
on the Rate of  $S_N2$  Reactions

Alkyl Halide	Relative Rate
$\text{CH}_3\text{CH}_2-$	1
$\text{CH}_3\text{CH}_2\text{CH}_2-$	0.4
$(\text{CH}_3)_2\text{CHCH}_2-$	0.03
$(\text{CH}_3)_3\text{CCH}_2-$	0.00001

FIGURE 8.8  $S_N2$  attack at two conformations of *n*-propyl compounds.FIGURE 8.9  $S_N2$  reaction at isobutyl systems

## Chap. 8

Reactions of  
Alkyl HalidesFIGURE 8.10  $S_N2$  reaction at neopentyl compounds.

Neopentyl halides are particularly interesting because there is no conform in which a blocking methyl group can be avoided (Figure 8.10). Neopentyl h are essentially unreactive in  $S_N2$  reactions except under very drastic cond

Substitution of sites more remote than the  $\beta$ -carbon have little or no eff the case of  $S_N2$  reactions. For example, *n*-butyl and *n*-pentyl halides re essentially the same rate as *n*-propyl halides.

The type of steric interaction we have discussed here forces groups to away from each other. Such deformation often forces orbitals to overlap noncolinear fashion, which provides less effective bonding than colinear or (Figure 8.11). Overlap of orbitals that are not colinear gives bonds which be described as "bent." Such bent bonds are generally of higher energy th corresponding "straight bonds." These principles apply not only to the tra states for displacement reactions but to certain strained molecules as well (C 23).




FIGURE 8.11 Bent bonds have reduced orbital overlap.

In summary, the effect of the structure of the alkyl group on the rate of reaction is apparent in two ways.

1. Branching at the  $\alpha$ -carbon hinders reaction: rate order is methyl > secondary  $\gg$  tertiary.
2. Branching at the  $\beta$ -carbon hinders reaction: neopentyl compounds are particularly slow.

Displacements that proceed by the  $S_N2$  mechanism are most successful with primary compounds having no branches at the  $\beta$ -carbon. Yields are poor with secondary halides and with primary halides having branches at C-2. Neopentyl systems undergo the reaction only under very drastic conditions and tertiary halides do not react by this mechanism at all. When the rate of the  $S_N2$  reaction is slowed down by these structural effects, alternative side reactions become competitive. With tertiary halides, and to an important degree with secondary and highly branched primary halides, the side reactions tend to dominate. These reactions are discussed in Sections 8.7 and 8.8.


## 8.6 Some Typical $S_N2$ Reactions

### A. Leaving Groups


### Sec. 8.6 Some Typical $S_N2$ Reactions

Alkyl chlorides, bromides, and iodides all react satisfactorily by the  $S_N2$  mechanism. The ease of reaction is dependent on the nature of the leaving group, alkyl iodides reacting most rapidly and alkyl chlorides most slowly. Alkyl fluorides are essentially unreactive by the  $S_N2$  mechanism. Since chlorine is much cheaper than bromine, alkyl chlorides are the least expensive alkyl halides. However, for laboratory uses where only small amounts of material are involved, alkyl bromides are commonly used because they are 50-100 times more reactive than the corresponding chlorides. Iodides are somewhat more reactive than bromides but are quite a bit more expensive, and this slightly increased reactivity does not justify their additional cost. In industrial processes, where massive amounts of materials are involved and cost is a prime consideration, alkyl chlorides are used almost exclusively.

The  $S_N2$  reaction is not restricted to alkyl halides. Any group that is the conjugate base of a strong acid can act as a leaving group. An example is bisulfate ion,  $\text{HSO}_4^-$ , which is the conjugate base of sulfuric acid,  $pK_a = 5$ . Dimethyl sulfate is an inexpensive commercial compound that reacts readily by the  $S_N2$  mechanism. The leaving group is the methylsulfate ion, which is similar in its base strength to bisulfate ion.




The chief disadvantage of dimethyl sulfate is its toxicity. It is water soluble and reacts readily with the nucleophilic groups in body tissues and fluids. Although dimethyl sulfate is the only sulfate in common use, alkyl sulfonates are often employed. Sulfonic acids,  $\text{RSO}_3\text{H}$ , are similar to sulfuric acid in acidity and the sulfonate ion,  $\text{RSO}_3^-$ , is an excellent leaving group. Alkyl benzenesulfonates, alkyl *p*-toluenesulfonates, and alkyl methanesulfonates are extremely useful substrates for  $S_N2$  reactions. These compounds are readily prepared from alcohols as described in Sections 11.7.D and 18.13.A.




Alkyl nitrates undergo reaction by the  $S_N2$  mechanism because nitric acid is a strong acid and nitrate ion is a weak base. However, alkyl nitrates are more prone to side reactions than the corresponding halides and, therefore, the yield of substitution product is lower. Consequently, nitrates are rarely used.

The facility with which a group can function as a leaving group in an  $S_N2$  reaction is related to its basicity. If a group is a weak base (that is, the conjugate base of a strong acid), it will generally be a "good" leaving group. This is readily

understood by considering the electronic structure of the transition state.




The negative charge which is introduced with the attacking nucleophile is distributed over several atoms in the transition state. The charge is borne most by the entering and leaving groups, as shown by the resonance structures of the transition state.





The leaving group has gained an appreciable amount of electron density going from reactant to transition state. The more this electron density or negative charge is stabilized, the lower is the energy of the transition state and the faster is the rate of reaction. The degree to which a group can accommodate a negative charge is also related to its affinity for a proton, its basicity. The acids  $\text{HCl}$ ,  $\text{HI}$ , and  $\text{H}_2\text{SO}_4$  are all strong acids because the anions  $\text{Cl}^-$ ,  $\text{Br}^-$ ,  $\text{I}^-$ , and  $\text{HSO}_4^-$  are stable anions. These anions are also good leaving groups in  $\text{S}_\text{N}2$  reactions.

$\text{HCN}$  is a weak acid ( $\text{p}K_\text{a} = 10$ ) and the displacement of cyanide is not observed.




Hydrazoic acid ( $\text{HN}_3$ ) and acetic acid ( $\text{CH}_3\text{CO}_2\text{H}$ ) are also weak acids ( $\text{p}K_\text{a}$  5.8 and 4.8, respectively). Correspondingly, azide ion and acetate ion are extremely poor leaving groups.




The reason that alkyl fluorides are ineffective substrates in the  $\text{S}_\text{N}2$  reaction is related to the relatively low acidity of  $\text{HF}$  ( $\text{p}K_\text{a} = 3$ ).

By comparing  $\text{S}_\text{N}2$  reactivity with relative acidity, we can understand the operation of acid catalysis in certain displacement processes. Alcohols undergo  $\text{S}_\text{N}2$  reactions because hydroxide ion is too basic (the  $\text{p}K_\text{a}$  of its conjugate acid,  $\text{H}_2\text{O}$ , is 15.7).



However, in the presence of a strong mineral acid, such as  $\text{HCl}$ ,  $\text{HBr}$ , or  $\text{HI}$ , the alcohol oxygen is protonated. An  $\text{S}_\text{N}2$  reaction can now occur because the leaving group is water, which is a much weaker base than  $\text{OH}^-$  (the conjugate acid of water,  $\text{H}_3\text{O}^+$ , has  $\text{p}K_\text{a} = -1.7$ ).




Note that the same principles of electron density and relative basicity are involved in this reaction, even though the leaving group is not an anion. This is an important reaction of alcohols and it will be developed more fully in Chapter 11.

## Sec. 8.6


Some Typical  
S<sub>N</sub>2 Reactions

## B. Solvents


A number of solvents may be used as reaction media for S<sub>N</sub>2 reactions. Ethanol and methanol are particularly useful because they are inexpensive, relatively inert, and dissolve many organic substrates and inorganic salts. Sometimes some additional water is added to increase the solubility of the inorganic salt used as the displacing agent. Some typical examples follow



*n*-Butyl bromide is refluxed with sodium methoxide in methanol for ½ hr. Water is added and the organic layer is separated, dried, and distilled to give methyl *n*-butyl ether.




Isopropyl bromide and sodium thiocyanate (NaSCN) are refluxed in 90% aqueous alcohol for 6 hr. The precipitated sodium bromide is filtered. The filtrate is diluted with water and extracted with ether. Distillation gives isopropyl thiocyanate, (CH<sub>3</sub>)<sub>2</sub>CHSCN, in 76-79% yield



Benzyl chloride, C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>Cl, is refluxed with sodium cyanide for 4 hr in aqueous alcohol. The sodium chloride is filtered, the solvent is distilled and the product benzyl cyanide, C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>CN, is distilled under vacuum to give a 80-90% yield. Sodium and potassium cyanides are highly toxic white solids that are very soluble in water and slightly soluble in ethanol. Organic cyanides may be hydrolyzed to carboxylic acids (Section 17.6).

The use of acetone as a solvent for halide exchange reactions was illustrated at the beginning of this chapter. Acetone is an example of a polar aprotic solvent, that is, a solvent without hydroxy groups but with a relatively high dipole moment and dielectric constant. Polar aprotic solvents are useful because of their ability to solvate ions and, thereby, to dissolve many salts. Other examples are acetonitrile, CH<sub>3</sub>CN; dimethylformamide, (CH<sub>3</sub>)<sub>2</sub>NCHO; dimethyl sulfoxide, (CH<sub>3</sub>)<sub>2</sub>SO; and hexamethylphosphoric triamide, [(CH<sub>3</sub>)<sub>3</sub>N]<sub>3</sub>PO.

Displacement reactions in polar aprotic solvents are frequently much faster than they are in hydroxylic solvents. For example, the reaction of methyl bromide with iodide ion is about 500 times faster in acetone than in methyl alcohol. An even more striking example is the reaction of chloride ion with methyl iodide, which is a million times faster in dimethylformamide than it is in methyl alcohol. The relative rate of reaction of azide ion with *n*-butyl bromide in various solvents is shown for comparison in Table 8.5. One important reason for this effect is hydrogen bonding. In a hydroxylic solvent, the OH groups can solvate anions by hydrogen bonding:




Hydrogen bonding will be discussed in detail in Section 11.3. For the present, suffice it to say that such bonding tends to tie up an anion and make it less reactive. Aprotic solvents have no hydrogens capable of hydrogen bonding. Hence, the anions are more free and have greater effective nucleophilicity.



TABLE 8.5  
Relative Rates for the Reaction  
 $N_3^- + n\text{-C}_4\text{H}_9\text{Br} \rightarrow n\text{-C}_4\text{H}_9\text{N}_3 + \text{Br}^-$

Solvent	Relative Rate
$\text{CH}_3\text{OH}$	1
$(\text{CH}_3)_2\text{SO}$ (DMSO)	1.300
$(\text{CH}_3)_2\text{NCHO}$ (DMF)	2.800
$\text{CH}_3\text{CN}$	5.000
$[(\text{CH}_3)_2\text{N}]_3\text{PO}$ (HMPT)	200.000


This type of solvation effect can even change the relative reactivities of nucleophilic groups. For example, the order of reactivity of halide ion reaction in water or in alcohols is



In acetone, the reactivities tend to be closer together and in dimethylformamide the order is even reversed! More basic anions tend to be hydrogen bonded more firmly and are less reactive in hydroxylic solvents. Consequently, nucleophilicity is not a simple and invariant property, but depends on the specific conditions, and solvents. We make frequent use of qualitative rather than quantitative generalizations, and it is still useful to know that iodide ion is more reactive than chloride ion, and that second-row elements are usually more reactive in displacement reaction than first-row elements. Even though they are generally more reactive in polar aprotic solvents than in aqueous solutions, the aqueous solvents are still frequently used in practice because they are inexpensive, convenient, and, for many reactions, serve perfectly adequately.

### C. Displacement Reactions at Atoms Other than Carbon

Displacement reactions are very common and may occur at atoms other than carbon. Many well-known inorganic reactions can be formulated as nucleophilic displacement reactions. One example is the formation of hypochlorite (bleach) by the reaction of hydroxide ion with chlorine. The reaction is considered as an  $\text{S}_\text{N}2$  reaction in which hydroxide ion displaces chloride from a chlorine molecule:







The preceding equation illustrates the use of arrows for "electron-pushing" device for keeping track of electron pairs when describing the bond reorganization that occurs during a reaction. An arrow symbolizes the flow of an electron pair in a reaction. The arrow begins at the electron pair in a reactant and points to the electron pair "goes" in the reaction. In the example given, an electron pair from the hydroxide ion attacks a chlorine and forms a new covalent bond. At the same time, the electron pair that had bonded the two chlorines together is released with the leaving chloride ion. It is conventional to use curved arrows.


Another example is the Raschig process for the synthesis of hydrazine, which involves two such displacement steps.

## Sec. 8.7


## E2 Elimination



Even ordinary acid-base reactions can be regarded as displacement reactions on hydrogen.





Such reactions are extremely rapid and we generally consider them only as facile equilibria. However, with special techniques, the rates of proton transfer reactions can be measured, even though the second-order rate constants are as high as  $10^{10} \text{ M}^{-1} \text{ sec}^{-1}$ . This is many orders of magnitude faster than the analogous second-order rate constants for displacements on carbon. The transition state for such a proton-transfer reaction may be formulated as



In such a transition state the 1s orbital of the hydrogen is partially bound to both the incoming and the leaving base.


Many reactions of atoms and radicals with molecules can be regarded as radical displacement reactions. Mechanistically, such displacements are classified as **bimolecular homolytic substitution** reactions,  $\text{S}_{\text{H}}2$ . We have already discussed such reactions in Chapter 5, but now they take their place in a family of related reactions.



## 8.7

## E2 Elimination

One of the side reactions that occurs in varying degree in displacement reactions is the elimination of the elements of HX to product an alkene.




Under appropriate conditions, this reaction can be the principal reaction and becomes a method for preparing alkenes. Accordingly, it is discussed in more detail in Section 12.5.A. For the present, it suffices to know that this reaction occurs by attack of a base on a hydrogen with concomitant formation of a C=C double bond and breaking of the C—X bond to form halide ion.

Mechanistically, the reaction is classified as bimolecular elimination, or E2. Since

## Sec. 18.5

## Synthesis


## 4. Transesterification (Section 18.9.B).



## 5. Acyl halides and alcohols (Section 18.9.B).




## 6. Acid anhydrides and alcohols (Section 18.9.B)




## Amides


## 1. Pyrolysis of ammonium carboxylates (Section 17.7.C)




## 2. Acyl halides and ammonia or amines (Section 18.9.C)






## 3. Acid anhydrides and ammonia or amines (Section 18.9.C)



## 4. Partial hydrolysis of nitriles (Section 18.9.A)



## Acyl Halides

1. Carboxylic acids with  $\text{SOCl}_2$  (Section 17.7.C).2. Carboxylic acids with  $\text{PCl}_5$  (Section 17.7.C).3. Carboxylic acids with  $\text{PBr}_3$  (Section 17.7.C)



A service of the National Library of Medicine  
and the National Institutes of Health

www.pubmed.gov

My NCBI [?] [Sign In] [Register]

All Databases  
PubMed

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

for camptothecin\* and cancer polychemotherapy

Go

Clear

Save Search

☒ Limits

☐ Preview/Index

☐ History

☐ Clipboard

☐ Details

Limits: Publication Date to 2002

Display: Summary

Show: 50

Sort by

Send to

Entrez  
Toolbox

Version

Free PubMed

View

FAQ

Help

Noteworthy  
Alerts

Med Services

PubMed Database

PubMed Database

PubMed Citation Matcher

PubMed Citation Matcher

PubMed Queries

PubMed Queries

NCBI

Related Resources

PubMed Documents

PubMed Mobile

PubMed Catalog

PubMed Gateway

PubMed NET

PubMed Health

PubMed Alerts

PubMed Alerts

PubMed Central

All: 649 Review: 212

Items 1 - 50 of 212

Page 1

of 5 Next

1: Wilkinson K.

Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut

**Irinotecan hydrochloride**  
Clin J Oncol Nurs. 2001 Jul-Aug;5(4):179-80, 184. Review. No abstract available.  
PMID: 12690622 [PubMed - indexed for MEDLINE]

2: Bleickardt E, Aramis A, Righ R, Blum K, McKee A, Tura H, Zeltman D, Burrows B, Davies M, Murren R.

Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut

**Phase I dose escalation trial of weekly docetaxel plus irinotecan in patients with advanced cancer.**  
Cancer Biol Ther. 2002 Nov-Dec;1(6):646-51. Review.  
PMID: 12642688 [PubMed - indexed for MEDLINE]

3: Hejblum V.

Related Articles, Compound via MeSH, Substance via MeSH, Cited in PMC, Books, LinkOut

**Gemcitabine in the treatment of advanced pancreatic cancer: a comparative analysis of randomized trials.**  
Semin Oncol. 2002 Dec;29(6 Suppl 20):9-16. Review.  
PMID: 12577228 [PubMed - indexed for MEDLINE]

4: Hochster H.

Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut

**The role of pemetrexed in the treatment of colorectal cancer.**  
Semin Oncol. 2002 Dec;29(6 Suppl 18):54-6. Review.  
PMID: 12571812 [PubMed - indexed for MEDLINE]

5: Rothenberg M.

Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut

**Current status of capecitabine in the treatment of colorectal cancer.**  
Oncology (Williston Park). 2002 Dec;16(12 Suppl No 14):16-22. Review.  
PMID: 12520635 [PubMed - indexed for MEDLINE]

6: Hobbay TJ, Goldberg RM.

Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut

**Perspectives on the role of sequential or combination chemotherapy for first-line and salvage therapy in advanced colorectal cancer.**  
Clin Colorectal Cancer. 2002 Nov;2(3):161-9. Review.  
PMID: 12482332 [PubMed - indexed for MEDLINE]

7: Douillard Y, Eckardt J, Souchon G.

Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut

**Challenging the platinum combinations in the chemotherapy of NSCLC.**  
Lung Cancer. 2002 Dec;38 Suppl 4:21-8. Review.  
PMID: 12480191 [PubMed - indexed for MEDLINE]

8: Asami T, Kokawa A, Sugimoto K, Tanaka N, Shirai K, Morimoto M, Saito T, Tanaka K, Imada T.

Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut

**[A case of AFP-producing gastric cancer with multiple liver metastases responding to CPT-11 and cisplatin combination chemotherapy]**  
Gan To Kagaku Ryoho. 2002 Nov;29(11):1985-8. Review. Japanese.  
PMID: 12465401 [PubMed - indexed for MEDLINE]


























9: Wasserman E, Sutherland W, Chikoye F.

Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut

**Irinotecan plus oxaliplatin: a promising combination for advanced colorectal cancer.**  
Clin Colorectal Cancer. 2001 Nov;1(3):149-53. Review.  
PMID: 12450427 [PubMed - indexed for MEDLINE]

10: Hobbay TJ, Frickman C.

Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut

-  Adjuvant therapy of colon cancer: a review.  
Clin Colorectal Cancer. 2002 Feb;1(4):230-6. Review.  
PMID: 12450421 [PubMed - indexed for MEDLINE]
-  11: Levyet C, Andre T, Arzu P. [Related Articles](#), [Compound via MeSH](#), [Substance via MeSH](#), [Books](#), [LinkOut](#)  
 [Gemcitabine-based combinations in inoperable pancreatic cancers]  
Bull Cancer. 2002 Aug;89 Spec No S96-101. Review. French.  
PMID: 12449038 [PubMed - indexed for MEDLINE]
-  12: Rousier P, Milly E. [Related Articles](#), [Compound via MeSH](#), [Substance via MeSH](#), [Books](#), [LinkOut](#)  
 Review of the role of CPT-11 in the treatment of colorectal cancer  
Clin Colorectal Cancer. 2001 Aug;1(2):87-94. Review.  
PMID: 12445366 [PubMed - indexed for MEDLINE]
-  13: Fisher MD. [Related Articles](#), [Compound via MeSH](#), [Substance via MeSH](#), [Books](#), [LinkOut](#)  
 Irinotecan/5-FU/leucovorin, oxaliplatin/5-FU/leucovorin, and oxaliplatin/irinotecan are each effective in the treatment of 5-FU-resistant advanced colorectal cancer.  
Clin Colorectal Cancer. 2001 Aug;1(2):85-6. Review. No abstract available.  
PMID: 12445365 [PubMed - indexed for MEDLINE]
-  14: Fisher MD. [Related Articles](#), [Compound via MeSH](#), [Substance via MeSH](#), [Books](#), [LinkOut](#)  
 Recommendation for irinotecan, 5-Fluorouracil, and leucovorin as first-line therapy for colorectal cancer.  
Clin Colorectal Cancer. 2001 Aug;1(2):82-4. Review. No abstract available.  
PMID: 12445364 [PubMed - indexed for MEDLINE]
-  15: Budman DR. [Related Articles](#), [Compound via MeSH](#), [Substance via MeSH](#), [Books](#), [LinkOut](#)  
 The search for enhanced therapeutic index in breast cancer: targeting the tumor and modulation of enzyme expression.  
Cancer Invest. 2002;20 Suppl 2:38-44. Review. No abstract available.  
PMID: 12442348 [PubMed - indexed for MEDLINE]
-  16: Takai N, Utsunomiya H, Kawano Y, Nishi K, Nishihara H, Miyakawa I. [Related Articles](#), [Compound via MeSH](#), [Substance via MeSH](#), [Books](#), [LinkOut](#)  
 Complete response to radiation therapy in a patient with chemotherapy-resistant ovarian clear cell adenocarcinoma.  
Arch Gynecol Obstet. 2002 Dec;267(2):98-100. Review.  
PMID: 12439556 [PubMed - indexed for MEDLINE]
-  17: Marx G, Harper P. [Related Articles](#), [Compound via MeSH](#), [Substance via MeSH](#), [Books](#), [LinkOut](#)  
 Non-platinum gemcitabine combinations in non-small cell lung cancer.  
Lung Cancer. 2002 Nov;38 Suppl 2:S51-4. Review.  
PMID: 12431830 [PubMed - indexed for MEDLINE]
-  18: Needle MN. [Related Articles](#), [Compound via MeSH](#), [Substance via MeSH](#), [Books](#), [LinkOut](#)  
 Safety experience with IMC-C225, an anti-epidermal growth factor receptor antibody.  
Semin Oncol. 2002 Oct;29(5 Suppl 14):55-60. Review.  
PMID: 12422314 [PubMed - indexed for MEDLINE]
-  19: Schmidt HJ. [Related Articles](#), [Compound via MeSH](#), [Substance via MeSH](#), [Books](#), [LinkOut](#)  
 The role of oxaliplatin in the treatment of advanced metastatic colorectal cancer: prospects and future directions.  
Semin Oncol. 2002 Oct;29(5 Suppl 15):34-9. Review.  
PMID: 12422306 [PubMed - indexed for MEDLINE]
-  20: Von Hoff DD, Reilly D. [Related Articles](#), [Compound via MeSH](#), [Substance via MeSH](#), [Books](#), [LinkOut](#)  
 New drugs for patients with pancreatic cancer.  
Curr Opin Oncol. 2002 Nov;14(6):621-7. Review.  
PMID: 12409652 [PubMed - indexed for MEDLINE]
-  21: Kelly K. [Related Articles](#), [Compound via MeSH](#), [Substance via MeSH](#), [Books](#), [LinkOut](#)  
 Current role of irinotecan in the treatment of non-small-cell lung cancer  
Oncology (Williston Park). 2002 Sep;16(9):1153-62, 1165; discussion 1165-6 passim. Review.  
PMID: 12380945 [PubMed - indexed for MEDLINE]
-  22: Sandler A. [Related Articles](#), [Compound via MeSH](#), [Substance via MeSH](#), [Books](#), [LinkOut](#)  
 Irinotecan plus cisplatin in small-cell lung cancer.  
Oncology (Williston Park). 2002 Sep;16(9 Suppl 9):39-43. Review.  
PMID: 12375800 [PubMed - indexed for MEDLINE]

- 23: Kwong MS, Bleickardt E, Murren JR  
 Camptothecin and taxane regimens for small-cell lung cancer.  
 Oncology (Williston Park). 2002 Sep;16(9 Suppl 9):33-8. Review.  
 PMID: 12375799 [PubMed - indexed for MEDLINE]
- 24: Uchino Y Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 [Randomized controlled trials in Japan--lung cancer]  
 Gan To Kagaku Ryoho. 2002 Sep;29(9):1516-21. Review. Japanese.  
 PMID: 12355937 [PubMed - indexed for MEDLINE]
- 25: Bunn PA Jr Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 Chemotherapy for advanced non-small-cell lung cancer: who, what, when, why?  
 J Clin Oncol. 2002 Sep 15;20(18 Suppl):23S-33S. Review.  
 PMID: 12235221 [PubMed - indexed for MEDLINE]
- 26: Bertin D Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 Targeting vascular endothelial growth factor in colorectal cancer.  
 Oncology (Williston Park). 2002 Aug;16(8 Suppl 7):13-5. Review.  
 PMID: 12199627 [PubMed - indexed for MEDLINE]
- 27: Green MR Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 Perspectives and opportunities: Docetaxel in the current and future treatment of non-small cell lung cancer.  
 Semin Oncol. 2002 Jun;29(3 Suppl 12):17-21. Review.  
 PMID: 12170447 [PubMed - indexed for MEDLINE]
- 28: Ferr A, Sanchez JM, Russell R, Taron M, Martinez F, Guillot M, Manzana JL, Mergel M, Ramadas A, Abad A Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 Phase I study of weekly CPT-11 (irinotecan)/docetaxel in patients with advanced solid tumors.  
 Lung Cancer. 2002 Aug;37(2):213-8. Review.  
 PMID: 12140145 [PubMed - indexed for MEDLINE]
- 29: Feyer G, Lianan B, Kraft D, Descoe L, Toller-Lennir V Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 Therapeutic advances in the management of metastatic colorectal cancer.  
 Expert Rev Anticancer Ther. 2001 Aug;1(2):236-46. Review.  
 PMID: 12113029 [PubMed - indexed for MEDLINE]
- 30: Masuda N, Fukuoka M Related Articles, Compound via MeSH, Substance via MeSH, Cited in Books, Books, LinkOut  
 Irinotecan in the treatment of small cell lung cancer.  
 Expert Rev Anticancer Ther. 2001 Aug;1(2):187-95. Review.  
 PMID: 12113024 [PubMed - indexed for MEDLINE]
- 31: Minsky BJ Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 Combined-modality therapy for rectal cancer using irinotecan  
 Oncology (Williston Park). 2002 May;16(5 Suppl 5):35-8. Review.  
 PMID: 12109805 [PubMed - indexed for MEDLINE]
- 32: Verschnagen CE Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 Irinotecan for the treatment of cervical cancer.  
 Oncology (Williston Park). 2002 May;16(5 Suppl 5):32-4. Review.  
 PMID: 12109804 [PubMed - indexed for MEDLINE]
- 33: Ikon DL, Minsky B, Kelsen D Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 Irinotecan, cisplatin, and radiation in esophageal cancer.  
 Oncology (Williston Park). 2002 May;16(5 Suppl 5):11-5. Review.  
 PMID: 12109799 [PubMed - indexed for MEDLINE]
- 34: Yoh K, Kubota K Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 [New treatment strategy for stage III non-small-cell lung cancer]  
 Nippon Rinsho. 2002 May;60 Suppl 5:485-9. Review. Japanese. No abstract available.  
 PMID: 12101719 [PubMed - indexed for MEDLINE]
- 35: Nokihara H, Qhe Y Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 [Recent progress in the treatment of ED small cell lung cancer]  
 Nippon Rinsho. 2002 May;60 Suppl 5:480-4. Review. Japanese. No abstract available.  
 PMID: 12101718 [PubMed - indexed for MEDLINE]

- 36: Yokoyama A. Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 [Irinotecan]  
 Nippon Rinsho. 2002 May;60 Suppl 5:339-42. Review. Japanese. No abstract available.  
 PMID: 12101684 [PubMed - indexed for MEDLINE]
- 37: Ohe Y. Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 Treatment-related death from chemotherapy and thoracic radiotherapy for advanced cancer.  
 Panminerva Med. 2002 Sep;44(3):205-12. Review.  
 PMID: 12094134 [PubMed - indexed for MEDLINE]
- 38: Kozuch P, Penyk M, Druckner HW. Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 A comprehensive update on the use of chemotherapy for metastatic pancreatic adenocarcinoma.  
 Hematol Oncol Clin North Am. 2002 Feb;16(1):123-38. Review.  
 PMID: 12063823 [PubMed - indexed for MEDLINE]
- 39: Tsuboi M, Kato H. Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 [Current status of the chemotherapy for lung cancer]  
 Gan To Kagaku Ryoho. 2002 May;29(5):684-94. Review. Japanese.  
 PMID: 12040671 [PubMed - indexed for MEDLINE]
- 40: Sandler A. Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 Irinotecan therapy for small-cell lung cancer.  
 Oncology (Williston Park). 2002 Apr;16(4):419-25, 428, 433, discussion 433-4, 437-8. Review.  
 PMID: 12017533 [PubMed - indexed for MEDLINE]
- 41: Kerr DJ. Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 Capecitabine/irinotecan combination regimens in colorectal cancer.  
 Oncology (Williston Park). 2002 Apr;16(4 Suppl 3):27-9. Review.  
 PMID: 12014865 [PubMed - indexed for MEDLINE]
- 42: Geyvindarajah R. Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 Irinotecan/thalidomide in metastatic colorectal cancer.  
 Oncology (Williston Park). 2002 Apr;16(4 Suppl 3):23-6. Review.  
 PMID: 12014864 [PubMed - indexed for MEDLINE]
- 43: Crino L. Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 [Role of irinotecan in the treatment of small cell carcinoma]  
 Tumori. 2001 Nov-Dec;87(6):A35-7. Review. Italian. No abstract available.  
 PMID: 11995707 [PubMed - indexed for MEDLINE]
- 44: Barone C. Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 [Irinotecan in the treatment of stomach cancer]  
 Tumori. 2001 Nov-Dec;87(6):A33-5. Review. Italian. No abstract available.  
 PMID: 11995706 [PubMed - indexed for MEDLINE]
- 45: Falcone A. Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 [Colorectal cancer, irinotecan in combination with newer drugs]  
 Tumori. 2001 Nov-Dec;87(6):A31-3. Review. Italian. No abstract available.  
 PMID: 11995705 [PubMed - indexed for MEDLINE]
- 46: Ando A, Shih CH. Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 [Chemotherapy for the patients with esophageal cancer]  
 Nippon Geka Gakkai Zasshi. 2002 Apr;103(4):359-63. Review. Japanese.  
 PMID: 11993225 [PubMed - indexed for MEDLINE]
- 47: Lloyd Jones M, Hummel S, Rana-Jack N, Cote B, Seymour M. Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 A rapid and systematic review of the evidence for the clinical effectiveness and cost-effectiveness of irinotecan, oxaliplatin and raltitrexed for the treatment of advanced colorectal cancer.  
 Health Technol Assess. 2001;5(25):1-128. Review. No abstract available.  
 PMID: 11990245 [PubMed - indexed for MEDLINE]
- 48: Ilson DH. Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut  
 New developments in the treatment of esophageal cancer.  
 Curr Oncol Rep. 2002 May;4(3):213-21. Review.  
 PMID: 11937011 [PubMed - indexed for MEDLINE]
- Mayradsis D. Related Articles, Compound via MeSH, Substance via MeSH, Books, LinkOut

49:



[New therapeutic approaches in non-small-cell lung carcinoma (NSCLC), stages IV]  
Rev Pneumol Clin. 2001 Nov;57(5 Pt 2):S42-5. Review. French. No abstract available.  
PMID: 11924243 [PubMed - indexed for MEDLINE]

50:



Hollen KD, Saliz LB.

[Related Articles](#), [Compound via MeSH](#), [Substance via MeSH](#), [Books](#), [LinkOut](#)

New therapies, new directions: advances in the systemic treatment of metastatic colorectal cancer.  
Lancet Oncol. 2001 May;2(5):290-7. Review.  
PMID: 11905784 [PubMed - indexed for MEDLINE]

Items 1 - 50 of 212

Page 1

of 5 [Next](#)Display **Summary**

Show

50



Sort by



Send to



[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Mar 10 2016 04:11:02



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**